

* FORCING

Garvin Melles*
Hebrew University of Jerusalem

February 1, 2008

Abstract

Let M be a transitive model of ZFC and let $\mathbf{B}$ be a M -complete Boolean algebra in M . (Possibly a proper class.) We define a generalized notion of forcing with such Boolean algebras, $*$ forcing. We prove that

1. If $\mathbf{G}$ is a $*$ forcing complete ultrafilter on $\mathbf{B}$, then $M[\mathbf{G}] \models ZFC$.
2. Let $H \subseteq M$. If there is a least transitive model N such that $H \in M$, $Ord^M = Ord^N$, and $N \models ZFC$, then we denote N by $M[H]$. We show that all models of ZFC of the form $M[H]$ are $*$ forcing extensions of M . As an immediate corollary we get that $L[0^\#]$ is a $*$ forcing extension of L .

1 Introduction

In this paper we introduce a generalization of forcing, $*$ forcing. If M is a transitive model of ZFC and $\mathbf{B}$ is a M -complete Boolean algebra in M , then a $*$ forcing extension of M is a transitive class of the form $M[\mathbf{G}]$ where $\mathbf{G}$ is an M -complete ultrafilter on $\mathbf{B}$. If $\mathbf{B}$ is a set Boolean algebra, then $*$ forcing and forcing coincide, so the interesting case is when $\mathbf{B}$ is a proper class. One indication of the importance of $*$ forcing is given by the fact that $L[0^\#]$ is a $*$ forcing extension of L . (In fact we show much more, as in the abstract.) In what follows we review Boolean valued models, and show how Boolean valued models on names for sets lead to pre- $*$ forcing,

*Would like to thank Ehud Hrushovski for supporting him with funds from NSF Grant DMS 8959511

* forcing, and forcing with Boolean values for formulas. We prove that if $M[\mathbf{G}]$ is a * forcing extension of M and $\mathbf{G}$ is * forcing complete, then $M[\mathbf{G}] \models ZFC$. In the final section we show that if $M[H]$ is the smallest transitive model of ZFC such that $Ord^M = Ord^{M[H]}$ and $H \in M$, then there is a * forcing complete ultrafilter $\mathbf{H}$ on the Boolean algebra $\mathbf{B}_{form}$ (a proper class in M) such that $M[H] = M[\mathbf{H}]$.

2 Boolean Valued Models

Definition 2.1 A Boolean valued model $M(B)$ (of the language of set theory) is a triple (U, I, E) where U is the universe and I and E are binary functions with values in some Boolean algebra. Keeping with tradition we write $I(x, y)$ as $\|x = y\|$ and $E(x, y)$ as $\|x \in y\|$. I and E must satisfy the following laws:

1. $\|x = x\| = 1$
2. $\|x = y\| = \|y = x\|$
3. $\|x = y\| \bullet \|y = z\| \leq \|x = z\|$
4. $\|x \in y\| \bullet \|v = x\| \bullet \|w = y\| \leq \|v \in w\|$

For each formula and $a_1, \dots, a_n$ from U we define the Boolean value of as follows:

1. $\|\neg\psi(a_1, \dots, a_n)\| = -\|\psi(a_1, \dots, a_n)\|$
2. $\|\varphi \wedge \psi(a_1, \dots, a_n)\| = \|\varphi(a_1, \dots, a_n)\| \bullet \|\psi(a_1, \dots, a_n)\|$
3. $\|\exists x \varphi(x, a_1, \dots, a_n)\| = \sum_{a \in M(B)} \|\varphi(a, a_1, \dots, a_n)\|$

If $M(B)$ is a Boolean valued model (or even if $\|\cdot\|$ is defined only for the atomic formulas) we can with any ultrafilter F on B construct a model $M(B)/F$, as follows: Define an equivalence relation over $M(B)$ by letting

$$x \equiv y \quad \text{iff} \quad \|x = y\| \in F$$

Define the relation E on the equivalence classes by

$$[x] E [y] \quad \text{iff} \quad \|x \in y\| \in F$$

Let $M(B)/F = (M(B)/\equiv, E)$.

Lemma 2.2 *If $M(B)$ is a Boolean valued model and F is an ultrafilter on B such that for every formula $\varphi = \exists x\psi(x)$ and $a_1, \dots, a_n$ from the universe of $M(B)$, if*

$$||\exists x\psi(x, a_1, \dots, a_n)|| \in F$$

implies there is an a in the universe of $M(B)$ such that

$$||\psi(a, a_1, \dots, a_n)|| \in F$$

then for every formula θ , and $b_1, \dots, b_m$ from the universe of $M(B)$,

$$M(B)/F \models \theta([b_1], \dots, [b_m]) \text{ iff } ||\theta(b_1, \dots, b_m)|| \in F$$

PROOF Like the proof of Lemma 18.1 in [Jech].

3 * Forcing and * Forcing Complete Ultrafilters

Definition 3.1 *If M is a transitive model of ZFC and $\mathbf{B}$ is a class Boolean algebra in M then we define the Boolean valued names for M , $M^{\mathbf{B}}$ as*

$$\bigcup_{\alpha \in \text{Ord}^M} M^{B_\alpha}$$

where $B_\alpha = \{b \in \mathbf{B} \mid \text{rank}(b) \leq \alpha\}$.

Definition 3.2 *Let M be a transitive model of ZFC and let $\mathbf{N}$ be a class in M . We call $\mathbf{N}$ in the rest of this definition the class of names. Let P be a class of unary relation symbols. We define the class of formulas $\text{Form}(M, \mathbf{N}, P)$ as the smallest class $\mathbf{C}$ such that*

1. *Each symbol in P is in $\mathbf{C}$, together with each symbol of P with a name substituted for the variable*
2. *$x = y$ and $x \in y$ are in $\mathbf{C}$, together with substitutions of variables in $x = y$ and $x \in y$ by names*
3. *If $\varphi(x_0, x_1, \dots, x_n, A)$ is in $\mathbf{C}$ then so are*

$$\neg\varphi(x_0, x_1, \dots, x_n, A)$$

$$\exists x_0\varphi(x_0, x_1, \dots, x_n, A)$$

4. Let I be a set. If for each $i \in I$, $\varphi(x_0, x_1, \dots, x_n, A_i) \in \mathbf{C}$, then

$$\bigwedge_{i \in I} \varphi(x_0, x_1, \dots, x_n, A_i) \in \mathbf{C}$$

$$\bigvee_{i \in I} \varphi(x_0, x_1, \dots, x_n, A_i) \in \mathbf{C}$$

Let $\text{Sent}(M, \mathbf{N}, P)$ be the elements of $\text{Form}(M, \mathbf{N}, P)$ without free variables. Let $\text{Sent}(M, \mathbf{N}, P)(q.f.)$ be the sentences in $\text{Sent}(M, M^{\mathbf{N}}, P)$ without quantifiers and similarly for $\text{Form}(M, \mathbf{N}, P)(q.f.)$.

Definition 3.3 A pre- $\ast$ forcing notion is a quadruple $(M, M^{\mathbf{B}}, \|\cdot\|^\ast, \mathbf{G})$ such that

1. M is a transitive model of ZFC
2. $M^{\mathbf{B}}$ is the class of Boolean valued names for M
3. $\|\cdot\|^\ast$ is a function with domain the elements of $\text{Sent}(M, M^{\mathbf{B}})(q.f.)$ such that
 - (a) $\|x = y\|^\ast = 0$ if $x \neq y$ and $\|x = y\|^\ast = 1$ if $x = y$
 - (b) $\|x \in y\|^\ast = b$ if $x \in \text{dom } y$ and $y(x) = b$ and 0 otherwise
 - (c) If I is a set and for each $i \in I$, A_i is a set of names, then
 - i. $\|\neg\psi(A_0)\|^\ast = -\|\psi(A_0)\|^\ast$
 - ii. $\|\bigvee_{i \in I} \varphi(A_i)\|^\ast = \sum_{i \in I} \|\varphi(A_i)\|^\ast$
 - iii. $\|\bigwedge_{i \in I} \varphi(A_i)\|^\ast = \prod_{i \in I} \|\varphi(A_i)\|^\ast$
4. $\mathbf{G}$ is a M -complete ultrafilter on $\mathbf{B}$, i.e., it has the property that if $X \subseteq \mathbf{B}$ and

$$\sum X \in \mathbf{G}$$

then there is an $x \in X$ such that $x \in \mathbf{G}$.

We denote $(M^{\mathbf{B}}, \|\cdot\|^\ast)$ by $M(\mathbf{B})$.

Theorem 3.4 If $(M, M^{\mathbf{B}}, \|\cdot\|^\ast, \mathbf{G})$ is a pre- $\ast$ forcing notion, then for every $\varphi(A) \in \text{Sent}(M, M^{\mathbf{B}})(q.f.)$,

$$M(\mathbf{B})/\mathbf{G} \models \varphi(A) \quad \text{iff} \quad \|\varphi(A)\|^\ast \in \mathbf{G}$$

PROOF By induction on the complexity of φ and the definition of pre- $\ast$ forcing notion.

So we have a notion of a condition forcing a formula, at least for quantifier free elements of $Sent(M, M^{\mathbf{B}})$. The disadvantage with pre- $\ast$ forcing is that the model $M(\mathbf{B})/\mathbf{G}$ is not in general a model of extensionality. But the function $\| \cdot \|^\ast$ induces a function $\| \cdot \|$ so that if we denote the $\ast$ Boolean valued model $(M, M^{\mathbf{B}}, \| \cdot \|)$ by $M^{\mathbf{B}}$, then extensionality holds in $M^{\mathbf{B}}/\mathbf{G}$. ($(M^{\mathbf{B}}, \| \cdot \|)$ is just the usual notion of Boolean valued model for forcing.)

Definition 3.5 *If $(M, M^{\mathbf{B}}, \| \cdot \|^\ast, \mathbf{G})$ is a pre- $\ast$ forcing notion then we define by induction on $\Gamma(\rho(x), \rho(y))$ where $\rho(x)$ is the rank of x and Γ is the canonical well ordering of $Ord \times Ord$,*

$$\|x \in y\| = \sum_{t \in \text{dom } y} \|x = t\| \bullet \|t \in y\|^\ast$$

$$\|x = y\| = \prod_{t \in \text{dom } x} (-\|t \in x\|^\ast + \|t \in y\|) \bullet \prod_{t \in \text{dom } y} (-\|t \in y\|^\ast + \|t \in x\|)$$

We call $(M, M^{\mathbf{B}}, \| \cdot \|, \mathbf{G})$ a $\ast$ forcing notion.

Theorem 3.6 *If $(M, M^{\mathbf{B}}, \| \cdot \|^\ast, \mathbf{G})$ is a pre- $\ast$ forcing notion then $\| \cdot \|$ and $M^{\mathbf{B}}/\mathbf{G}$ with respect to $\| \cdot \|$ are defined and*

$$M^{\mathbf{B}}/\mathbf{G} \cong M[\mathbf{G}]$$

by the isomorphism $h[x] = i_{\mathbf{G}}(x)$ where

$$M[\mathbf{G}] = \{i_{\mathbf{G}}(x) \mid x \in M^{\mathbf{B}}\}$$

with $\in$ on $M[\mathbf{G}]$ interpreted as the real $\in$ relation and $i_{\mathbf{G}}$ is defined in the usual manner.

PROOF See lemmata 18.2, 18.3, 18.9 and exercise 18.6 of [Jech].

Theorem 3.7 *(The forcing theorem for $\ast$ forcing) If $\varphi(A) \in Sent(M, M^{\mathbf{B}})(q.f.)$, then*

$$M[\mathbf{G}] \models \varphi(A) \quad \text{iff} \quad \|\varphi(A)\| \in \mathbf{G}$$

PROOF For every sentence $\varphi(A) \in \text{Sent}(M, M^{\mathbf{B}})$ there is a corresponding statement $\varphi'(A')$ such that $\|\varphi(A)\| = \|\varphi'(A')\|^*$ and

$$M^{\mathbf{B}}/\mathbf{G} \models \varphi(A) \leftrightarrow M(\mathbf{B})/\mathbf{G} \models \varphi'(A')$$

Furthermore, $\varphi(A) \in \text{Sent}(M, M^{\mathbf{B}})(q.f.) \leftrightarrow \varphi'(A') \in \text{Sent}(M, M^{\mathbf{B}})(q.f.)$
 If $(M, M^{\mathbf{B}}, \|\cdot\|^*, \mathbf{G})$ is a pre- $*$ forcing notion and if $\varphi'(A') \in \text{Sent}(M, M^{\mathbf{B}})(q.f.)$ we know that

$$M(\mathbf{B})/\mathbf{G} \models \varphi'(A') \leftrightarrow \|\varphi'(A')\|^* \in \mathbf{G}$$

This lifts to

$$M^{\mathbf{B}}/\mathbf{G} \models \varphi(A) \leftrightarrow \|\varphi(A)\| \in \mathbf{G}$$

Since $M[\mathbf{G}]$ and $M^{\mathbf{B}}/\mathbf{G}$ are isomorphic we get the statement in the theorem.

Definition 3.8 Let $(M, M^{\mathbf{B}}, \|\cdot\|, \mathbf{G})$ be a $*$ forcing notion. Suppose that for every first order formula $\varphi(x_0, x_1, \dots, x_n)$ and set of names $\dot{a}_1, \dots, \dot{a}_n$ there is a set $B_{\varphi(x_0, \dot{a}_1, \dots, \dot{a}_n)}$ of names such that for every $C \supset B_{\varphi(x_0, \dot{a}_1, \dots, \dot{a}_n)}$,

$$\sum_{\dot{b} \in B_{\varphi(x_0, \dot{a}_1, \dots, \dot{a}_n)}} \|\varphi(\dot{b}, \dot{a}_1, \dots, \dot{a}_n)\| = \sum_{\dot{c} \in C} \|\varphi(\dot{c}, \dot{a}_1, \dots, \dot{a}_n)\|$$

where the definition of $\|\cdot\|$ is extended to all first order formulas by induction on their complexity as follows:

1. $\|\neg\varphi(\dot{a}_1, \dots, \dot{a}_n)\| = -\|\varphi(\dot{a}_1, \dots, \dot{a}_n)\|$
2. $\|\varphi(\dot{a}_1, \dots, \dot{a}_n) \wedge \psi(\dot{a}_1, \dots, \dot{a}_n)\| = \|\varphi(\dot{a}_1, \dots, \dot{a}_n)\| \bullet \|\psi(\dot{a}_1, \dots, \dot{a}_n)\|$
- 3.

$$\|\exists x_0 \varphi(x_0, \dot{a}_1, \dots, \dot{a}_n)\| = \sum_{\dot{b} \in B_{\varphi(x_0, \dot{a}_1, \dots, \dot{a}_n)}} \|\varphi(\dot{b}, \dot{a}_1, \dots, \dot{a}_n)\|$$

Then we say $(M, M^{\mathbf{B}}, \|\cdot\|, \mathbf{G})$ is a forcing notion with Boolean values for formulas.

Theorem 3.9 If $(M, M^{\mathbf{B}}, \|\cdot\|, \mathbf{G})$ is a forcing notion with Boolean values for formulas, then

$$M[\mathbf{G}] \models \text{ZFC} - P$$

PROOF First we show that the forcing theorem holds for all first order formulas by induction on the complexity of the formulas, and then the usual proof that a forcing extension (using a set Boolean algebra) satisfies *ZFC* for all the axioms except power set goes through.

Definition 3.10 Let S be a class. A formula $\varphi(v_1, \dots, v_n)$ is said to be a formula of set theory enriched by S if it is a first order formula built from the usual atomic formulas of set theory together with atomic formulas of the form $s(x)$ for $s \in S$ where the interpretation of $s(x)$ is $x \in s$.

Definition 3.11 If $\alpha < \text{Ord}^M$, let $M_\alpha^{\mathbf{B}}$ be the names of rank $\leq \alpha$, and let $M_\alpha^{B_\alpha}$ be the elements of M^{B_α} of rank $\leq \alpha$.

Definition 3.12 Let $\alpha < \text{Ord}^M$ and let $(M, M^{\mathbf{B}}, \|\cdot\|, \mathbf{G})$ be a $*$ forcing notion. Let $\varphi(x_1, \dots, x_n)$ be a formula and let $\dot{a}_1, \dots, \dot{a}_n$ be names in $M_\alpha^{B_\alpha}$. By induction on the complexity of φ we define

$$\|\varphi(\dot{a}_1, \dots, \dot{a}_n)\|_\alpha$$

as follows:

1. $\|\varphi(\dot{a}_1, \dots, \dot{a}_n)\|_\alpha = \|\varphi(\dot{a}_1, \dots, \dot{a}_n)\|$ if φ is quantifier free.
2. $\|\varphi(\dot{a}_1, \dots, \dot{a}_n) \wedge \psi(\dot{a}_1, \dots, \dot{a}_n)\|_\alpha = \|\varphi(\dot{a}_1, \dots, \dot{a}_n)\|_\alpha \bullet \|\psi(\dot{a}_1, \dots, \dot{a}_n)\|_\alpha$
3. $\|\neg\varphi(\dot{a}_1, \dots, \dot{a}_n)\|_\alpha = -\|\varphi(\dot{a}_1, \dots, \dot{a}_n)\|_\alpha$
4. $\|\exists x \varphi(\dot{a}_1, \dots, \dot{a}_n)\|_\alpha = \sum_{b \in M_\alpha^{B_\alpha}} \|\psi(b, \dot{a}_1, \dots, \dot{a}_n)\|_\alpha$

Definition 3.13 Let $\alpha < \text{Ord}^M$ and let $(M, M^{\mathbf{B}}, \|\cdot\|, \mathbf{G})$ be a $*$ forcing notion. Let $\varphi(x_1, \dots, x_n)$ be a formula. We define by induction on the complexity of φ the meaning of $\varphi(x_1, \dots, x_n)$ reflects in $M_\alpha^{B_\alpha}$.

1. If $\varphi(x_1, \dots, x_n)$ is quantifier free then $\varphi(x_1, \dots, x_n)$ reflects in every $M_\alpha^{B_\alpha}$
2. $\varphi(x_1, \dots, x_n)$ reflects in $M_\alpha^{B_\alpha}$ iff $\neg\varphi(x_1, \dots, x_n)$ reflects in $M_\alpha^{B_\alpha}$
3. $\varphi(x_1, \dots, x_n) \wedge \psi(x_1, \dots, x_n)$ reflects in $M_\alpha^{B_\alpha}$ iff $\varphi(x_1, \dots, x_n)$ and $\psi(x_1, \dots, x_n)$ reflect in $M_\alpha^{B_\alpha}$

4. If $\varphi(x_1, \dots, x_n) = \exists x_0 \psi(x_0, x_1, \dots, x_n)$ then $\varphi(x_1, \dots, x_n)$ reflects in $M_\alpha^{B_\alpha}$ iff

- (a) $\psi(x_0, x_1, \dots, x_n)$ reflects in $M_\alpha^{B_\alpha}$
- (b) $\forall \beta > \alpha \exists \gamma > \beta$ such that $\psi(x_0, x_1, \dots, x_n)$ reflects in $M_\gamma^{B_\gamma}$ and if $b \in M_\gamma^{B_\gamma}$ and $\|\psi(b, \dot{a}_1, \dots, \dot{a}_n)\|_\gamma \in \mathbf{G}$, for some $\dot{a}_1, \dots, \dot{a}_n$ from $M_\alpha^{B_\alpha}$ then there is an $a \in M_\alpha^{B_\alpha}$ such that

$$\|\psi(a, \dot{a}_1, \dots, \dot{a}_n)\|_\alpha \in \mathbf{G}$$

Definition 3.14 If $(M, M^{\mathbf{B}}, \|\cdot\|, \mathbf{G})$ is a $*$ forcing notion then we say that $\mathbf{G}$ is $*$ forcing complete if

- 1. For every $\alpha < \text{Ord}^M$, there is an ordinal β such that if $\dot{a}$ is a name with domain $M_\alpha^{B_\alpha}$, then for some name $\dot{b}$ in $M_\beta^{B_\beta}$

$$\|\dot{a} = \dot{b}\| \in \mathbf{G}$$

- 2. For every formula of set theory $\varphi(x_1, \dots, x_n)$ and for every $\alpha < \text{Ord}^M$, there is an ordinal $\beta > \alpha$ such that $\varphi(x_1, \dots, x_n)$ reflects in $M_\beta^{B_\beta}$

Theorem 3.15 If $(M, M^{\mathbf{B}}, \|\cdot\|, \mathbf{G})$ is a $*$ forcing notion and $\mathbf{G}$ is $*$ forcing complete, then

$$M[\mathbf{G}] \models ZFC$$

PROOF The only nontrivial axioms to prove are choice, union, power set, replacement and separation. Separation follows from replacement and from separation it is not hard to prove union.

Replacement First we prove by induction on the complexity of φ that if φ reflects in $M_\alpha^{B_\alpha}$ then $\forall \dot{a}_1, \dots, \dot{a}_n \in M_\alpha^{B_\alpha}$

$$M[\mathbf{G}] \models \varphi(i_{\mathbf{G}}(\dot{a}_1), \dots, i_{\mathbf{G}}(\dot{a}_n)) \leftrightarrow i_{\mathbf{G}}[M_\alpha^{B_\alpha}] \models \varphi(i_{\mathbf{G}}(\dot{a}_1), \dots, i_{\mathbf{G}}(\dot{a}_n))$$

$$\leftrightarrow \|\varphi(\dot{a}_1, \dots, \dot{a}_n)\|_\alpha \in \mathbf{G}$$

(The second $\leftrightarrow$ follows from the definition of $\|\cdot\|_\alpha$.) If φ is quantifier free then it follows immediately. The only nontrivial step is if $\varphi = \exists x \psi(x)$.

Suppose $M[\mathbf{G}] \models \varphi(i_{\mathbf{G}}(\dot{a}_1), \dots, i_{\mathbf{G}}(\dot{a}_n))$. Then for some $\beta < Ord^M$ there is a $\dot{b} \in M_{\beta}^{B_{\beta}}$ such that

$$M[\mathbf{G}] \models \psi(i_{\mathbf{G}}(\dot{b}), i_{\mathbf{G}}(\dot{a}_1), \dots, i_{\mathbf{G}}(\dot{a}_n))$$

Since $\mathbf{G}$ is $\ast$ forcing complete there is a $\gamma > \beta$ such that $\psi(x_0, x_1, \dots, x_n)$ reflects in $M_{\gamma}^{B_{\gamma}}$ which implies by the induction hypothesis that

$$i_{\mathbf{G}}[M_{\gamma}^{B_{\gamma}}] \models \psi(i_{\mathbf{G}}(\dot{b}), i_{\mathbf{G}}(\dot{a}_1), \dots, i_{\mathbf{G}}(\dot{a}_n))$$

and $\|\psi(\dot{b}, \dot{a}_1, \dots, \dot{a}_n)\|_{\gamma} \in \mathbf{G}$. Since $\mathbf{G}$ is $\ast$ forcing complete and $\varphi(x_1, \dots, x_n)$ reflects in $M_{\alpha}^{B_{\alpha}}$, $\psi(x_0, x_1, \dots, x_n)$ reflects in $M_{\alpha}^{B_{\alpha}}$ and for some $\dot{a} \in M_{\alpha}^{B_{\alpha}}$,

$$\|\psi(\dot{a}, \dot{a}_1, \dots, \dot{a}_n)\|_{\alpha} \in \mathbf{G}$$

and $i_{\mathbf{G}}[M_{\alpha}^{B_{\alpha}}] \models \psi(i_{\mathbf{G}}(\dot{a}), i_{\mathbf{G}}(\dot{a}_1), \dots, i_{\mathbf{G}}(\dot{a}_n))$ which implies

$$\|\varphi(\dot{a}_1, \dots, \dot{a}_n)\|_{\alpha} \in \mathbf{G}$$

and $i_{\mathbf{G}}[M_{\alpha}^{B_{\alpha}}] \models \varphi(i_{\mathbf{G}}(\dot{a}_1), \dots, i_{\mathbf{G}}(\dot{a}_n))$. Now if $\|\varphi(\dot{a}_1, \dots, \dot{a}_n)\|_{\alpha} \in \mathbf{G}$ then we know that since $\mathbf{G}$ is M -complete that for some $a \in M_{\alpha}^{B_{\alpha}}$,

$$\|\psi(\dot{a}, \dot{a}_1, \dots, \dot{a}_n)\|_{\alpha} \in \mathbf{G}$$

Now by the induction hypothesis we get

$$M[\mathbf{G}] \models \psi(i_{\mathbf{G}}(\dot{a}), i_{\mathbf{G}}(\dot{a}_1), \dots, i_{\mathbf{G}}(\dot{a}_n))$$

i.e., $M[\mathbf{G}] \models \varphi(i_{\mathbf{G}}(\dot{a}_1), \dots, i_{\mathbf{G}}(\dot{a}_n))$. Now let $F(x) = y$ be a function defined by the formula $\varphi(x, y, i_{\mathbf{G}}(\dot{a}_1), \dots, i_{\mathbf{G}}(\dot{a}_n))$ with the $\dot{a}_i$ from $M_{\alpha}^{B_{\alpha}}$. Let $\dot{X}$ be a set and let $i_{\mathbf{G}}(\dot{X}) = X$. Let $\beta > \alpha$ such that $\varphi(x, y, \dot{a}_1, \dots, \dot{a}_n)$ reflects in $M_{\beta}^{B_{\beta}}$ and $\dot{X} \in M_{\beta}^{B_{\beta}}$. Let $\dot{Y}$ be the name with domain $M_{\beta}^{B_{\beta}}$ such that $\forall \dot{a} \in M_{\beta}^{B_{\beta}}$,

$$\dot{Y}(\dot{a}) = \sum_{x \in \text{dom } \dot{X}} \|\varphi(x, \dot{a}, \dot{a}_1, \dots, \dot{a}_n)\|_{\beta} \bullet \|x \in \dot{X}\|$$

Then $F[X] = i_{\mathbf{G}}(\dot{Y})$.

Power Set By point 1. of the definition of $\ast$ forcing complete and by separation we know that for each $\alpha < Ord^M$ that $P(i_{\mathbf{G}}(M_{\alpha}^{B_{\alpha}}))$ exists.

Now applying separation and the fact that if $X \in M[\mathbf{G}]$, then for some $\alpha < Ord^M$, $X \subseteq i_{\mathbf{G}}[M_{\alpha}^{B_{\alpha}}]$ we get the power set axiom holds in $M[\mathbf{G}]$.

Choice If X is a name a set in $M[\mathbf{G}]$ and f is the name with domain $= \{(\check{x}, x)^{\mathbf{B}} \mid x \in dom X\}$ such that $f((\check{x}, x)^{\mathbf{B}})$ is uniformly 1, then $i_{\mathbf{G}}(f)$ is a function from $dom X$ onto $i_{\mathbf{G}}(X)$. Since $dom X$ is well ordered (as it is in M) $i_{\mathbf{G}}(f)$ induces a well order on $i_{\mathbf{G}}(X)$.

Corollary 3.16 *Let T be a theory extending ZFC . If $(M, M^{\mathbf{B}}, || \cdot ||, \mathbf{G})$ is a $*$ forcing notion and $\mathbf{G}$ is $*$ forcing complete, then*

$$M[\mathbf{G}] \models T$$

iff $\forall \varphi \in T, \exists \alpha < Ord^M$ such that $M_{\alpha}^{B_{\alpha}}$ reflects φ and $||\varphi||_{\alpha} \in \mathbf{G}$.

We would like necessary and sufficient conditions that $M[\mathbf{G}] \models ZFC$ but the proof does not seem to go in the other direction unless $\mathbf{G}$ is a class in $M[\mathbf{G}]$. But if $\mathbf{G}$ is a class in $M[\mathbf{G}]$ then the $\{i_{\mathbf{G}}(M_{\alpha}^{B_{\alpha}}) \mid \alpha < Ord^M\}$ form a cumulative hierarchy in $M[\mathbf{G}]$ and we can prove:

Theorem 3.17 *If $M[\mathbf{G}]$ is a $*$ forcing extension of M , $\mathbf{G}$ is a class in $M[\mathbf{G}]$ and $M[\mathbf{G}] \models ZFC$ then $\mathbf{G}$ is $*$ forcing complete.*

PROOF If $\mathbf{G}$ is a class in $M[\mathbf{G}]$ then the $\{i_{\mathbf{G}}(M_{\alpha}^{B_{\alpha}}) \mid \alpha < Ord^M\}$ form a cumulative hierarchy in $M[\mathbf{G}]$ so that for every formula φ and for every ordinal $\alpha < Ord^M$, there is a $\beta > \alpha$ such that φ reflects (in the usual sense) in $i_{\mathbf{G}}(M_{\beta}^{B_{\beta}})$. Now we can show by induction on the complexity of φ that φ reflects in $M_{\beta}^{B_{\beta}}$. Since the power set axiom holds in $M[\mathbf{G}]$ we know that condition 1. in the definition of $*$ forcing complete must also hold, otherwise for some α the collection of subsets of $i_{\mathbf{G}}(M_{\alpha}^{B_{\alpha}})$ form a class in $M[\mathbf{G}]$.

If we consider models of set theory with an addition predicate $G(x)$ meant as an interpretation for $\mathbf{G}$ we can get necessary and sufficient conditions that $M[\mathbf{G}] \models ZFC_{G(x)}$ with the interpretation of $G(x)$ being $\mathbf{G}$ where we define $ZFC_{G(x)}$ as ZFC except that we allow in the separation and replacement schemata the formulas to be enriched by $G(x)$. So for the rest of this section if we write $M[\mathbf{G}]$ we mean the model in the language $(=, \in, G(x))$ with the interpretation of $G(x)$ being $\mathbf{G}$. If $(M, M^{\mathbf{B}}, || \cdot ||, \mathbf{G})$ is a $*$ forcing notion then we can extend the definition of $|| \cdot ||$ so that

$(M, M^{\mathbf{B}}, \Vdash)$ is a $\ast$ Boolean valued model in the language $(=, \in, G(x))$ so that if $\dot{a}$ is a name then

$$i_{\mathbf{G}}(\dot{a}) \in \mathbf{G} \quad \text{iff} \quad M[\mathbf{G}] \models G(i_{\mathbf{G}}(\dot{a})) \quad \text{iff} \quad \Vdash G(\dot{a}) \in \mathbf{G}$$

If $b \in \mathbf{B}$, let

$$G(\check{b}) = b$$

Otherwise, if $\dot{a}$ is a name in $M_{\alpha}^{\mathbf{B}}$, let

$$\Vdash G(\dot{a}) = \sum_{b \in B_{\alpha}} \Vdash \dot{a} = \check{b} \bullet \Vdash G(\check{b})$$

Now we can extend the definition of $\Vdash$ to sentences in $Sent(M, M^{\mathbf{B}}, \{G(x)\})(q.f.)$ in the natural way.

Definition 3.18 *If $(M, M^{\mathbf{B}}, \Vdash, \mathbf{G})$ is a $\ast$ forcing notion then we say that $\mathbf{G}$ is $\ast$ forcing complete relative to formulas enriched by $G(x)$ if*

1. *For every $\alpha < Ord^M$, there is an ordinal β such that if $\dot{a}$ is a name with domain $M_{\alpha}^{B_{\alpha}}$, then for some name $\dot{b}$ in $M_{\beta}^{B_{\beta}}$*

$$\Vdash \dot{a} = \dot{b} \in \mathbf{G}$$

2. *For every formula of set theory enriched by $G(x)$, $\varphi(x_1, \dots, x_n)$ and for every $\alpha < Ord^M$, there is an ordinal $\beta > \alpha$ such that $\varphi(x_1, \dots, x_n)$ reflects in $M_{\beta}^{B_{\beta}}$*

Theorem 3.19 *If $(M, M^{\mathbf{B}}, \Vdash, \mathbf{G})$ is a $\ast$ forcing notion then $M[\mathbf{G}] \models ZFC_{G(x)}$ iff $\mathbf{G}$ is $\ast$ forcing complete relative to formulas of set theory enriched by $G(x)$.*

PROOF Similar.

4 $\ast$ Forcing and models of ZFC of the form $M[H]$

Definition 4.1 *If U is a transitive set, then $Def_A(U)$ is the set of all subsets of U definable by formulas of set theory enriched by $A(x)$ with quantifiers restricted to U and parameters from U .*

Definition 4.2 If M is a transitive model of ZFC and $H \subset M$, then if there is a least transitive model N of ZFC such that $H \in M$ and $\text{Ord}^M = \text{Ord}^N$, then we denote N by $M[H]$.

Theorem 4.3 Let M be a transitive model of ZFC. If $H \subseteq M$ and $M[H]$ exists, then

$$M[H] = \bigcup_{\alpha \in \text{Ord}^M} L_\alpha(M \cup \{H\})$$

where

1. $L_0(M \cup \{H\}) = \emptyset$
2. $L_\alpha(M \cup \{H\}) = \bigcup_{\beta < \alpha} L_\beta(M \cup \{H\})$ if α is a limit
3. $L_{\alpha+1}(M \cup \{H\}) = \bigcup_{A \in M \cup \{H\}} \text{Def}_A L_\alpha(M \cup \{H\})$

PROOF See [Jech] exercise 15.13.

The above theorem induces a natural notion of **names** and of interpretation function i_H .

Definition 4.4 The class **names** is $\bigcup_{\alpha \in \text{Ord}^M} \text{names}_\alpha$ where we define names_α by induction on the ordinals in M as follows:

1. $\text{names}_0 = \emptyset$
2. $\text{names}_\alpha = \bigcup_{\beta < \alpha} \text{names}_\beta$ if α is a limit
3. $\tilde{y} \in \text{names}_{\alpha+1}$ iff $\tilde{y}$ is a function with domain names_α and $\tilde{y}(\tilde{z}) = \varphi^\alpha(\tilde{z}, \tilde{z}_1, \dots, \tilde{z}_n)$ where $\varphi(v_0, v_1, \dots, v_n)$ is a formula of set theory enriched by $M \cup \{H\}$ and the $\tilde{z}_i$ are in names_α

Definition 4.5 If $M[H]$ exists we define by induction on $\alpha < \text{Ord}^M$, i_H as follows:

1. $i_H(\emptyset) = \emptyset$

2. If $\tilde{y}$ is a name in $names_{\alpha+1}$ with the form as in the definition of **names** above, then

$$i_H(\tilde{y}) = \{i_H(\tilde{z}) \mid i_H[names_\alpha] \models \varphi(i_H(\tilde{z}), i_H(\tilde{z}_1), \dots, i_H(\tilde{z}_n))\}$$

$$\text{where } i_H[names_\alpha] = \{i_H(\tilde{z}) \mid \tilde{z} \in names_\alpha\} = L_\alpha(M \cup \{H\})$$

Definition 4.6 $\mathbf{B}_{form}$ is the class $Sent(M, \mathbf{names}, M \cup \{H\})(q.f.)$ made into a M -complete Boolean algebra by equating sentences logically equivalent in propositional logic. We have

1. $0 = [\emptyset = \emptyset \wedge \emptyset \neq \emptyset]$
2. $1 = [\emptyset = \emptyset \vee \emptyset \neq \emptyset]$
3. $-\lceil\varphi\rceil = \lceil\neg\varphi\rceil$
4. If $\{A_i \mid i \in I\}$ is a set of sets of names, then

$$(a) \prod \{[\varphi(A_i)] \mid i \in I\} = [\bigwedge_{i \in I} \varphi_i(A_i)]$$

$$(b) \sum \{[\varphi(A_i)] \mid i \in I\} = [\bigvee_{i \in I} \varphi(A_i)]$$

Definition 4.7 We define for every $\alpha < Ord^M$ a function Asn_α and a class function $\mathbf{I}$ such that

1. $dom Asn_\alpha =$ symbols of the form $\varphi^\alpha(x_1, \dots, x_m, \tilde{a}_1, \dots, \tilde{a}_n)$ where $\tilde{a}_1, \dots, \tilde{a}_n$ are in $names_\alpha$
2. $range Asn_\alpha \subseteq Form(M, \mathbf{names}, M \cup \{H\})(q.f.)$
3. $dom \mathbf{I} = \mathbf{names}$
4. $range \mathbf{I} \subseteq M^{\mathbf{B}_{form}}$

We define Asn_α by induction on the complexity of φ .

1. If φ is quantifier free, then

$$Asn_\alpha(\varphi^\alpha) = [\varphi]$$

2. $Asn_\alpha(\neg\varphi^\alpha) = -[Asn_\alpha(\varphi^\alpha)]$

$$3. \text{Asn}_\alpha(\varphi^\alpha \wedge \psi^\alpha) = [\text{Asn}_\alpha(\varphi^\alpha)] \bullet [\text{Asn}_\alpha(\psi^\alpha)]$$

$$4. \text{Asn}_\alpha(\exists x \psi(x)^\alpha) = [\bigvee_{\tilde{a} \in \text{names}_\alpha} \text{Asn}_\alpha(\psi^\alpha(\tilde{a}))]$$

We let $\mathbf{I} = \bigcup_{\alpha \in \text{Ord}^M} I_\alpha$ where we define I_α by induction on α as follows:

Let $I_0(\emptyset) = \emptyset$. Suppose I_α has been defined. Now if $\tilde{y} \in \text{name}_{\alpha+1}$ with elements in the range of $\tilde{y}$ of the form $\varphi^\alpha(\tilde{a}, \tilde{a}_1, \dots, \tilde{a}_n)$ then $I_{\alpha+1}(\tilde{y})$ has domain

$$\{I_\alpha(\tilde{a}) \mid \tilde{a} \in \text{names}_\alpha\}$$

and

$$I_{\alpha+1}(\tilde{y})(I_\alpha(\tilde{a})) = \text{Asn}_\alpha(\varphi(\tilde{a}, \tilde{a}_1, \dots, \tilde{a}_n))$$

Theorem 4.8 *Every model of the form $M[H]$ is a $*$ forcing extension of M via $\mathbf{B}_{\text{form}}$ by a $*$ forcing complete ultrafilter.*

PROOF Let $\mathbf{H} = \{[\varphi(A)] \mid \varphi(A) \in \text{Sent}(M, \mathbf{names}, M \cup \{H\})(q.f.) \wedge M[H] \models \varphi(i_H[A])\}$. We leave to the reader the proof that $\mathbf{H}$ is an ultrafilter and $\mathbf{H}$ is M -complete. Since $\mathbf{H}$ is definable inside $M[H]$ (If $\varphi(A) \in \text{Sent}(M, \mathbf{names}, M \cup \{H\})(q.f.)$ then $M[H] \models \varphi(i_H[A])$ iff $\exists \alpha (A \subseteq \text{names}_\alpha \wedge i_H[\text{names}_\alpha] \models \varphi(i_H[A]))$) we know that $M[\mathbf{H}] = i_{\mathbf{H}}[M^{\mathbf{B}_{\text{form}}}] \subseteq M[H]$. To prove the other direction we first prove by induction on the rank of $\tilde{y}$ that

$$i_H(\tilde{y}) = i_{\mathbf{H}}(\mathbf{I}(\tilde{y}))$$

For $\tilde{y} = \emptyset$ it follows from the definitions. So let $\alpha = \beta + 1$ and let $\tilde{y} \in \text{names}_\alpha - \text{names}_\beta$. Suppose $\tilde{y}(\tilde{a}) = \varphi^\alpha(\tilde{a}, \tilde{a}_1, \dots, \tilde{a}_n)$. Then

$$L_\alpha(M \cup \{H\}) \models \varphi(i_H(\tilde{a}), i_H(\tilde{a}_1), \dots, i_H(\tilde{a}_n)) \text{ iff}$$

$$L_\alpha(M \cup \{H\}) \models \text{Asn}_\alpha \varphi(i_H(\tilde{a}), i_H(\tilde{a}_1), \dots, i_H(\tilde{a}_n)) \text{ iff}$$

$$M[H] \models \text{Asn}_\alpha \varphi(i_H(\tilde{a}), i_H(\tilde{a}_1), \dots, i_H(\tilde{a}_n)) \text{ iff}$$

$$\text{Asn}_\alpha \varphi(\tilde{a}, \tilde{a}_1, \dots, \tilde{a}_n) \in \mathbf{H}$$

As a result,

$$\begin{aligned} i_H(\tilde{y}) &= \{i_H(\tilde{a}) \mid L_\alpha(M \cup \{H\}) \models \varphi(i_H(\tilde{a}), i_H(\tilde{a}_1), \dots, i_H(\tilde{a}_n))\} \\ &= \{i_H(\tilde{a}) \mid \text{Asn}_\alpha \varphi(\tilde{a}, \tilde{a}_1, \dots, \tilde{a}_n) \in \mathbf{H}\} \end{aligned}$$

$$\begin{aligned}
&= \{i_{\mathbf{H}}(\mathbf{I}(\tilde{a})) \mid \text{Asn}_\alpha \varphi(\tilde{a}, \tilde{a}_1, \dots, \tilde{a}_n) \in \mathbf{H}\} \\
&= i_{\mathbf{H}}(\mathbf{I}(\tilde{y}))
\end{aligned}$$

So we have $M[H] = i_H[\mathbf{names}] \subseteq i_{\mathbf{H}}[M^{\mathbf{B}_{form}}] = M[\mathbf{H}]$. (As $M[H] \models ZFC$ and $\mathbf{H}$ is a class in $M[H]$ we know $\mathbf{H}$ must be $\ast$ forcing complete.)

As a corollary, we get the following.

Corollary 4.9 $L[0^\#] = L[\mathbf{G}]$ where $\mathbf{G}$ is a $\ast$ forcing complete ultrafilter on $\mathbf{B}_{form}^L$.

REFERENCES

[Jech] T. Jech, *Set Theory*, Academic Press, 1978.